

Effective QCD Partition Function in Sectors with Non-Zero Topological Charge and Itzykson-Zuber Type Integral

Toshinao AKUZAWA*and Miki WADATI

*Department of Physics, Graduate School of Science, University of Tokyo,
Hongo 7-3-1, Bunkyo-ku, Tokyo, 113*

February 1, 2008

Abstract

It was conjectured by Jackson *et.al.* that the finite volume effective partition function of QCD with the topological charge $M - N$ coincides with the Itzykson-Zuber type integral for $M \times N$ rectangular matrices. In the present article we give a proof of this conjecture, in which the original Itzykson-Zuber integral is utilized.

1 Introduction

Let us start with quantum chromodynamics (QCD) with N flavors and a nonzero vacuum angle θ . Assume that the quarks have masses $m_1, m_2, \dots, m_N$. We introduce the quark-mass matrix $\mathbf{M}$ by

$$\mathbf{M} = \text{diag}(m_1, \dots, m_N) . \quad (1.1)$$

Using matrices γ_μ , $\mu = 1, 2, 3, 4$, defined by

$$\gamma_4 = i \text{diag}(1, 1, -1, -1) \quad (1.2)$$

and

$$\{\gamma_\mu, \gamma_\nu\} = -2\delta_{\mu\nu} , \quad (1.3)$$

the Lagrangian density in the Euclidean space is expressed as [1]

$$L = \frac{1}{4g^2} \sum_{\mu, \nu=1}^N \sum_{a=1}^{N^2-1} F_{\mu\nu}^a F_{\mu\nu}^a - i\theta\omega - i\bar{q} \sum_{\mu=1}^4 \gamma_\mu (\partial_\mu + iA_\mu) q + \bar{q}_R \mathbf{M} q_L + \bar{q}_L \mathbf{M}^\dagger q_R , \quad (1.4)$$

where

$$\omega = \frac{1}{32\pi^2} \text{tr} \epsilon^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta} . \quad (1.5)$$

We assume that the space-time constitutes the four-dimensional torus with the volume V . The winding number (or topological charge) ν of the gauge field, which is expressed as

$$\nu = \int_V d^4x \omega(x) , \quad (1.6)$$

is a topological invariant. The partition function $\mathcal{Z}(\theta)$ is written as

$$\mathcal{Z}(\theta) = \sum_{\nu=-\infty}^{\infty} e^{i\nu\theta} \mathcal{Z}_\nu , \quad (1.7)$$

where $\mathcal{Z}_\nu$ is the partition function in the sector with topological charge ν . We consider the region in which the volume V is large and

$$V\sigma\mathcal{S} \lesssim 1 , \quad (1.8)$$

*akuzawa@monet.phys.s.u-tokyo.ac.jp

where σ and $\mathcal{S}$ represent respectively the quark condensate and the quark mass scale. In the low-energy theory, the Goldstone bosons dominate the partition function. In the present case the Goldstone bosons are described by matrix fields $U(x)$ in $SU(N)$. The partition function takes the form

$$\mathcal{Z} = \int_{U(x) \in SU(N)} \delta U \exp \left\{ - \int_V d^4 x \left[\frac{f^2}{4} \text{tr}(\partial_\mu U^\dagger \partial_\mu U) - \sigma \text{Re}(\exp(i\theta/N) \text{tr} \mathbf{M} U^\dagger) + \dots \right] \right\}, \quad (1.9)$$

where f is the pion-decay constant and Re means the real part. By setting $U(x) = U_0 U_1(x)$ and ignoring the non-zero modes fluctuations associated with $U_1(x)$, we arrive at the effective QCD partition function in terms of the collective variable U_0 [1],

$$\mathcal{Z} = \int_{SU(N)} d\mu(U_0) \exp \left\{ V \sigma \text{Re} \left(e^{i\theta/N} \text{tr}(\mathbf{M} U_0^\dagger) \right) \right\}. \quad (1.10)$$

Hereafter we denote by μ the normalized measure with the two-sided invariance. From (1.10) it is deduced that $\mathcal{Z}_\nu$ is expressed as an integral over $U(N)$:

$$\mathcal{Z}_\nu = \int_{U(N)} d\mu(U) (\det U)^\nu \exp(V \sigma \text{Re} \text{tr} \mathbf{M} U). \quad (1.11)$$

In case quarks have equal masses, the $U(N)$ -integral in (1.11) is carried out in [1]. For different quark masses the explicit expression for $\mathcal{Z}_{\nu=0}$ is obtained in [2]. The key of the derivation in [2] is the Itzykson-Zuber type integral for complex matrices

$$\begin{aligned} & \int_{U(M)} d\mu(U) \int_{U(N)} d\mu(U') \exp(\text{Re} \text{tr}(U' \Psi U \Xi)) \\ &= \frac{2^{N(M-1)}}{N!} \prod_{j=0}^N (1+j)!(M-N+j)! \prod_{i=1}^N (x_i y_i)^{\frac{1}{2}} \frac{\det\{I_{M-N}(x_k y_l)\}_{k,l}}{\xi_0(x) \xi_0(y)}, \end{aligned} \quad (1.12)$$

where

$$\Xi = \begin{pmatrix} x_1 & 0 & \dots & 0 \\ 0 & x_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & x_N \\ \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 0 \end{pmatrix}, \quad (1.13)$$

$$\Psi = \begin{pmatrix} y_1 & 0 & \dots & 0 & \dots & 0 \\ 0 & y_2 & \dots & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & y_N & \dots & 0 \end{pmatrix}, \quad (1.14)$$

and $\xi_0(x)$ is defined by

$$\xi_0(x) = \prod_{i,j=1}^N (x_i^2 - x_j^2) \prod_{i=1}^N x_i^{M-N+\frac{1}{2}}. \quad (1.15)$$

There is a straightforward correspondence between $\mathcal{Z}_{\nu=0}$ and (1.12) for $M = N$. What was conjectured in [2] is that $\mathcal{Z}_\nu$ for general ν also corresponds to (1.12) for $M = N + \nu$. In the present article we shall give a proof to this conjecture. In our proof we make use of the original Itzykson-Zuber integral[3].

2 Correspondence between Itzykson-Zuber type Integral and Partition Function

The correspondence between $\mathcal{Z}_\nu$ and the integral (1.12) is easily certified. We assume that $M \geq N$ without losing any generality. We set

$$\Psi = v \mathcal{M}, \quad (2.1)$$

where $v = V\sigma$ is a real positive number and $\mathcal{M}$ is an $N \times M$ matrix such as

$$\mathcal{M} = \begin{pmatrix} m_1 & 0 & \dots & 0 & \dots & 0 \\ 0 & m_2 & \dots & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & m_N & \dots & 0 \end{pmatrix}. \quad (2.2)$$

Set also

$$\Xi = \mathbf{1}_{M,N} \equiv \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 0 \end{pmatrix}. \quad (2.3)$$

Then the Itzykson-Zuber type integral (1.12) becomes

$$G_{M,N}(\mathcal{M}) \stackrel{\text{def}}{=} \int_{U(M)} d\mu(U) \int_{U(N)} d\mu(U') \exp(\text{Re } v \text{tr}(U' \mathcal{M} U \mathbf{1}_{M,N})) . \quad (2.4)$$

In case $M = N$ the double integral in (2.4) reduces to a single integral:

$$G_{N,N}(\mathcal{M}) = \int_{U(N)} d\mu^M(U) \exp(\text{Re } v \text{tr}(\mathcal{M} U)) . \quad (2.5)$$

This is exactly the partition function $\mathcal{Z}_0$ in the sector with topological charge $\nu = 0$. Furthermore, by giving examples which suggest the existence of the correspondence for general ν , it was conjectured in [2] that

$$G_{M,N}(\mathcal{M}) = \text{constant} \times \mathcal{Z}_{M-N} . \quad (2.6)$$

3 Proof of the Conjecture

For definiteness sake we shall add a suffix which specifies the number of the flavors to the partition function in the sector with topological charge ν , that is,

$$\mathcal{Z}_\nu^N = \int_{U(N)} d\mu(U) (\det U)^\nu \exp(v \text{Re } \text{tr} MU) . \quad (3.1)$$

By properly choosing $S \in SU(N)$ and $\Lambda = \text{diag}(e^{ix_1}, e^{ix_2}, \dots, e^{ix_N})$, where $x_i \in \mathbb{R}$, the matrix $U \in U(N)$ is expressed as

$$U = S \Lambda S^\dagger . \quad (3.2)$$

The well-known transformation of the variables of the integrations [4] reads

$$\int_{U(N)} d\mu(U) = \frac{1}{N!(2\pi)^N} \int_{SU(N)} d\mu(S) \left(\prod_{i=1}^N \int_0^{2\pi} dx_i \right) \prod_{i < j} \sin^2 \frac{x_i - x_j}{2} . \quad (3.3)$$

In terms of S and Λ , (3.1) is transformed to

$$\mathcal{Z}_\nu^N = \frac{1}{N!(2\pi)^N} \prod_{i=1}^N \left(\int_0^{2\pi} dx_i e^{ix_i} \right) \prod_{i < j} \sin^2 \frac{x_i - x_j}{2} \int_{SU(N)} d\mu(S) \exp(v \text{tr} M S \Lambda S^\dagger) . \quad (3.4)$$

The expression (3.4) has a great advantage because the $SU(N)$ integration is readily carried out by using the original Itzykson-Zuber integral[3],

$$I(A, B; \beta) = \int_{SU(N)} d\mu(U) \exp[\beta \text{tr} A U B U^\dagger] \quad (3.5)$$

$$= \beta^{-N(N-1)/2} \prod_{n=1}^{N-1} n! \frac{\det \{ e^{\beta a_i b_j} \}_{1 \leq i, j \leq N}}{\Delta(a) \Delta(b)} , \quad (3.6)$$

where Δ is the Vandermonde determinant, for instance,

$$\Delta(a) = \prod_{1 \leq i < j \leq N} (a_i - a_j) . \quad (3.7)$$

Consequently there remain the integrations over N variables:

$$\mathcal{Z}_\nu^N = v^{-N(N-1)/2} \prod_{n=1}^{N-1} n! \prod_{i=1}^N \left(\int_0^{2\pi} dx_i e^{i\nu x_i} \right) \prod_{i < j} \sin^2 \frac{x_i - x_j}{2} \frac{\det \{e^{v m_j \cos x_i}\}_{1 \leq i, j \leq N}}{\Delta(\cos x) \Delta(m)} . \quad (3.8)$$

In the next place, let us examine $G_{M,N}(\mathcal{M})$. It can be shown that

$$G_{M,N}(\mathcal{M}) = G_{M,M}(\tilde{\mathcal{M}}) = \int d\mu^M(U) \exp \left(\text{Re } v \text{tr}(\tilde{\mathcal{M}}U) \right) , \quad (3.9)$$

where

$$\tilde{\mathcal{M}} = \text{diag}(m_1, m_2, \dots, m_N, 0, \dots, 0) . \quad (3.10)$$

For the right-hand-side of (3.9) a similar kind of decomposition as for $\mathcal{Z}_\nu^N$ is possible. As a result, $G_{M,N}(\mathcal{M})$ is rewritten as

$$G_{M,N}(\mathcal{M}) = \frac{v^{-M(M-1)/2}}{M!(2\pi)^M} \prod_{n=1}^{M-1} n! \prod_{i=1}^M \left(\int_0^{2\pi} dx_i \right) \prod_{i < j} \sin^2 \frac{x_i - x_j}{2} \frac{\det \{e^{v \tilde{m}_j \cos x_i}\}_{1 \leq i, j \leq M}}{\Delta(\cos x) \Delta(\tilde{m})} , \quad (3.11)$$

where

$$\tilde{m}_i = \begin{cases} m_i & (1 \leq i \leq N) \\ 0 & (N+1 \leq i \leq M) \end{cases} . \quad (3.12)$$

We shall prove the conjecture that

$$G_{M,N}(\mathcal{M}) = \text{constant} \times \mathcal{Z}_{M-N}^N , \quad (3.13)$$

by comparing (3.8) with (3.11).

We start from (3.11). The fraction

$$\frac{\det \{e^{v \tilde{m}_j \cos x_i}\}_{1 \leq i, j \leq M}}{\Delta(\cos x) \Delta(\tilde{m})} \quad (3.14)$$

in (3.11) is decomposed by using the Schur function $S_{\{\lambda\}}$ as

$$\begin{aligned} &= \prod_{i=1}^M \left(\sum_{k_i=0}^{\infty} \frac{v^{k_i}}{k_i!} \right) \frac{\prod_{j=1}^M \cos^{k_j} x_j \det \tilde{m}_j^{k_i}}{\Delta(\cos x) \Delta(\tilde{m})} \\ &= \sum_{k_M=0}^{\infty} \sum_{k_{M-1}(>k_M)}^{\infty} \cdots \sum_{k_1(>k_2)}^{\infty} \prod_{i=1}^M \left(\frac{v^{k_i}}{k_i!} \right) \frac{\det \cos^{k_i} x_j \det \tilde{m}_j^{k_i}}{\Delta(\cos x) \Delta(\tilde{m})} \\ &= \sum_{\lambda_M=0}^{\infty} \sum_{\lambda_{M-1}(\geq \lambda_M)}^{\infty} \cdots \sum_{\lambda_1(\geq \lambda_2)}^{\infty} \prod_{i=1}^M \left(\frac{v^{\lambda_i + M - i}}{(\lambda_i + M - i)!} \right) S_{\{\lambda\}}(\cos x) S_{\{\lambda\}}(\tilde{m}) . \end{aligned} \quad (3.15)$$

For our purpose, it is important to note that the Schur functions have the property[5],

$$\begin{aligned} S_{\{\lambda\}}(\tilde{m}) &= S_{\{\lambda\}}(m_1, \dots, m_N, 0, \dots, 0) \\ &= \begin{cases} S_{\{\lambda_1, \dots, \lambda_N\}}(m_1, \dots, m_N) & : \text{ if } \lambda_{N+1} = \lambda_{N+2} = \dots = \lambda_M = 0 \\ 0 & : \text{ otherwise} \end{cases} . \end{aligned} \quad (3.16)$$

It follows from (3.16) that

$$\begin{aligned} \frac{\det \{e^{v \tilde{m}_j \cos x_i}\}_{1 \leq i, j \leq M}}{\Delta(\cos x) \Delta(\tilde{m})} &= \prod_{i=1}^{M-N} \frac{v^{i-1}}{(i-1)!} \sum_{\lambda_N=0}^{\infty} \sum_{\lambda_{N-1}(\geq \lambda_N)}^{\infty} \cdots \sum_{\lambda_1(\geq \lambda_2)}^{\infty} \\ &\quad \times \prod_{i=1}^N \left(\frac{v^{\lambda_i + M - i}}{(\lambda_i + M - i)!} \right) S_{\{\lambda_1, \dots, \lambda_N, 0, \dots, 0\}}(\cos x) S_{\{\lambda_1, \dots, \lambda_N\}}(m) . \end{aligned} \quad (3.17)$$

Since the notational convention of the Schur function is rather inconvenient to the discussions below, we introduce a function $T_{\{k_1, \dots, k_L\}}^{(L)}$ defined by

$$T_{\{k_1, \dots, k_L\}}^{(L)}(x_1, \dots, x_L) = \frac{\det\{x_i^{k_j}\}_{1 \leq i, j \leq L}}{\Delta(x)}. \quad (3.18)$$

The Schur function and $T_{\{k_1, \dots, k_L\}}^{(L)}$ are related by

$$S_{\{\lambda_1, \dots, \lambda_L\}} = T_{\{k_1+L-1, k_2+L-2, \dots, k_{L-1}+1, k_L\}}^{(L)}. \quad (3.19)$$

In terms of $T_{\{k_1, \dots, k_L\}}^{(L)}(x_1, \dots, x_L)$, (3.17) is rewritten as

$$\begin{aligned} \frac{\det\{e^{v\tilde{m}_j \cos x_i}\}_{1 \leq i, j \leq M}}{\Delta(\cos x)\Delta(\tilde{m})} &= \prod_{i=1}^{M-N} \frac{v^{i-1}}{(i-1)!} \sum_{\lambda_N=0}^{\infty} \sum_{\lambda_{N-1}(\geq \lambda_N)}^{\infty} \cdots \sum_{\lambda_1(\geq \lambda_2)}^{\infty} \\ &\times \prod_{i=1}^N \left(\frac{v^{\lambda_i+M-i}}{(\lambda_i+M-i)!} \right) T_{\{\lambda_1+M-1, \dots, \lambda_N+M-N, M-N-1, \dots, 1, 0\}}^{(M)}(\cos x) T_{\{\lambda_1+N-1, \dots, \lambda_N\}}^{(N)}(m) \\ &= \prod_{i=1}^{M-N} \frac{v^{i-1}}{(i-1)!} \sum_{k_N=0}^{\infty} \sum_{k_{N-1}(\geq k_N)}^{\infty} \cdots \sum_{k_1(\geq k_2)}^{\infty} \\ &\times \prod_{i=1}^N \left(\frac{v^{k_i}}{k_i!} m_i^{N-M} \right) T_{\{k_1, \dots, k_N, M-N-1, \dots, 1, 0\}}^{(M)}(\cos x) T_{\{k_1, \dots, k_N\}}^{(N)}(m). \end{aligned} \quad (3.20)$$

Note that

$$T_{\{k_1, \dots, k_N, M-N-1, M-N-2, \dots, 1, 0\}}^{(M)}(\cos x) = \frac{1}{\Delta(\cos x)} \sum_{P \in \sigma_M} (-1)^P \prod_{i=1}^N \cos^{k_i} x_{P_i} \prod_{\alpha=1}^{M-N} \cos^{M-N-i} x_{P_{N+\alpha}}, \quad (3.21)$$

where σ_M is the permutation of $\{1, 2, \dots, M\}$. By introducing σ_M^N by

$$\sigma_M^N = \{P \in \sigma_M : P_{N+1} > P_{N+2} > \dots > P_M, P_1 > P_2 > \dots > P_N\}, \quad (3.22)$$

we may express the left hand side of (3.21) in terms of $T_{\{k_1, \dots, k_N\}}^{(N)}$ as

$$\begin{aligned} &T_{\{k_1, \dots, k_N, M-N-1, M-N-2, \dots, 1, 0\}}^{(M)}(\cos x) \\ &= \frac{1}{\Delta(\cos x)} \sum_{P \in \sigma_M^N} (-1)^P \det\{\cos^{k_i} x_{P_j}\}_{1 \leq i, j \leq N} \det\{\cos^{M-N-i} x_{P_{N+j}}\}_{1 \leq i, j \leq M-N} \\ &= \sum_{P \in \sigma_M^N} T_{\{k_1, \dots, k_N\}}^{(N)}(\cos x_{P_1}, \cos x_{P_2}, \dots, \cos x_{P_N}) \prod_{i=1}^N \prod_{j=N+1}^M \frac{1}{\cos x_{P_i} - \cos x_{P_j}}. \end{aligned} \quad (3.23)$$

Putting (3.11), (3.20) and (3.23) together we get

$$\begin{aligned} G_{M,N}(\mathcal{M}) &= \frac{v^{N(N-2M+1)/2}}{M!(2\pi)^M} \prod_{n=M-N}^{M-1} n! \sum_{k_N=0}^{\infty} \sum_{k_{N-1}(\geq k_N)}^{\infty} \cdots \sum_{k_1(\geq k_2)}^{\infty} \prod_{i=1}^N \left(\frac{v^{k_i}}{k_i!} m_i^{N-M} \right) T_{\{k_1, \dots, k_N\}}^{(N)}(m) \\ &\times \prod_{i=1}^M \left(\int_0^{2\pi} dx_i \right) \prod_{i < j}^M \sin^2 \frac{x_i - x_j}{2} \sum_{P \in \sigma_M^N} T_{\{k_1, \dots, k_N\}}^{(N)}(\cos x_{P_1}, \cos x_{P_2}, \dots, \cos x_{P_N}) \\ &\times \prod_{i=1}^N \prod_{\alpha=N+1}^M \frac{1}{\cos x_{P_i} - \cos x_{P_\alpha}} \\ &= \frac{v^{N(N-2M+1)/2}}{M!(2\pi)^M} M C_N \prod_{n=M-N}^{M-1} n! \sum_{k_1 \geq k_2 \geq \dots \geq k_N} \prod_{i=1}^N \left(\frac{v^{k_i}}{k_i!} m_i^{N-M} \right) T_{\{k_1, \dots, k_N\}}^{(N)}(m) \\ &\times \prod_{i=1}^N \left(\int_0^{2\pi} dx_i \right) \prod_{i < j}^N \sin^2 \frac{x_i - x_j}{2} T_{\{k_1, \dots, k_N\}}^{(N)}(\cos x_1, \cos x_2, \dots, \cos x_N) \\ &\times \prod_{i=N+1}^M \left(\int_{0+i\epsilon}^{2\pi+i\epsilon} dx_i \right) \prod_{N+1 \leq i < j \leq M} \sin^2 \frac{x_i - x_j}{2} \prod_{i=1}^N \prod_{\alpha=N+1}^M \frac{\sin^2 \frac{x_i - x_\alpha}{2}}{\cos x_i - \cos x_\alpha}. \end{aligned} \quad (3.24)$$

The infinitesimal positive number ϵ is introduced to avoid the “pseudo” poles at $x_i = -x_\alpha$. With a change of variables $y_i = e^{ix_i}$, the integration over $\{x_i : N+1 \leq i \leq M\}$ in the last expression in (3.24) is transformed to the contour integral as

$$2^{-(M-N)(M-1)} (-1)^{(M-N)(M-N-1)/2} \prod_{i=N+1}^M \left(-i \oint_{C_\epsilon} \frac{dy_i}{y_i} \right) \\ \times \prod_{N+1 \leq i < j \leq M} (y_i - y_j)(y_i^{-1} - y_j^{-1}) \prod_{i=1}^N \prod_{\alpha=N+1}^M \frac{y_\alpha - y_i}{y_\alpha y_i - 1} \quad (3.25)$$

where C_ϵ is a circle with radius $e^{-\epsilon}$ centered at the origin of the complex plane. Equation (3.25) is immediately integrated out:

$$= 2^{-(M-N)(M-1)} (2\pi)^{M-N} (-1)^{(M-N)(M+N-1)/2} (M-N)! \prod_{i=1}^N y_i^{M-N}. \quad (3.26)$$

Inserting (3.26) in (3.24) we find that

$$G_{M,N}(\mathcal{M}) = C_{(M,N)} \sum_{k_1 \geq k_2 \geq \dots \geq k_N} \prod_{i=1}^N \left(\frac{v^{k_i}}{k_i!} m_i^{N-M} \right) T_{\{k_1, \dots, k_N\}}^{(N)}(m) \\ \times \prod_{i=1}^N \left(\int_0^{2\pi} dx_i e^{i(M-N)x_i} \right) \prod_{i < j}^N \sin^2 \frac{x_i - x_j}{2} T_{\{k_1, \dots, k_N\}}^{(N)}(\cos x_1, \cos x_2, \dots, \cos x_N), \quad (3.27)$$

where

$$C_{(M,N)} = v^{N(N-2M+1)/2} 2^{-(M-N)(M-1)} (2\pi)^{M-N} (-1)^{(M-N)(M+N-1)/2} \frac{1}{N!} \prod_{n=M-N}^{M-1} n!. \quad (3.28)$$

The summation in (3.27) reads

$$\sum_{k_1 \geq k_2 \geq \dots \geq k_N} \prod_{i=1}^N \left(\frac{v^{k_i}}{k_i!} m_i^{N-M} \right) T_{\{k_1, \dots, k_N\}}^{(N)}(m) T_{\{k_1, \dots, k_N\}}^{(N)}(\cos x_1, \cos x_2, \dots, \cos x_N) \\ = \frac{1}{\Delta(\cos x) \Delta(m)} \sum_{k_1 \geq k_2 \geq \dots \geq k_N} \prod_{i=1}^N \left(\frac{v^{k_i}}{k_i!} m_i^{N-M} \right) \det \{m_i^{k_j}\}_{1 \leq i, j \leq N} \det \{\cos^{k_j} x_i\}_{1 \leq i, j \leq N} \\ = \frac{1}{\Delta(\cos x) \Delta(m)} \prod_{i=1}^N m_i^{N-M} \det \left\{ \exp(v m_i \cos x_j) \right\}_{1 \leq i, j \leq N}. \quad (3.29)$$

Finally by combining (3.27) and (3.29) we obtain

$$G_{M,N}(\mathcal{M}) = C_{(M,N)} \prod_{i=1}^N m_i^{N-M} \prod_{i=1}^N \left(\int_0^{2\pi} dx_i e^{i(M-N)x_i} \right) \\ \times \prod_{i < j}^N \sin^2 \frac{x_i - x_j}{2} \frac{1}{\Delta(\cos x) \Delta(m)} \det \left\{ \exp(v m_i \cos x_j) \right\}_{1 \leq i, j \leq N}, \quad (3.30)$$

which is proportional to (3.8) with $\nu = M - N$. This proves the conjecture (3.13).

4 Discussions

The partition function (1.11) with $\nu = 0$ has the two-sided $U(N)$ invariance. That makes it possible to carry out the integration. If $\nu \neq 0$, the partition function is invariant under $SU(N)$ but not invariant under $U(N)$. That is, $U(1)$ symmetry is broken. Despite an impression that the remaining $SU(N)$ symmetry is still large, it is not easy to treat the partition function analytically.

It is known [6] that the Itzykson-Zuber type integral for rectangular matrices is derived by considering the zero-curvature limit of the isotropic diffusion on the cosets $SU(M+N)/S(U(M) \times U(N))$ and $SU(M, N)/S(U(M) \times U(N))$. These cosets are classified as the Riemannian symmetric space of type AIII in the mathematical context[7]. We have proved in the present article the conjecture (3.13) that the partition functions in the sectors with non-zero topological charges equals the Itzykson-Zuber type integral for rectangular matrices up to the normalization. This correspondence makes it possible to apply a full knowledge of the symmetric spaces to the analysis of the sectors with $\nu \neq 0$.

References

- [1] H.Leutwyler and A.Smilga, Spectrum of Dirac operator and role of winding number in QCD. *Phys.Rev.D*, **46**(12), 5607 (1992).
- [2] A. Jackson, M. Şener, and J. Verbaarschot, Finite volume partition functions and Itzykson-Zuber integrals. *Phys.Lett.B*, **387**, 355 (1996).
- [3] C.Itzykson and J.-B.Zuber, The planar approximation. II. *J.Math.Phys.*, **21**, 411 (1980).
- [4] M.L.Mehta, *Random Matrices*. Academic Press, second edn. (1991).
- [5] I.G.Macdonald, *Symmetric Functions and Hall Polynomials*. Oxford University Press, Oxford, second edn. (1995).
- [6] T.Akuzawa and M.Wadati, Diffusions on symmetric spaces of type AIII and random matrix theories for rectangular matrices. *J.Phys.A*, **31**, 1713 (1998).
- [7] S.Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*. Academic Press, New York (1978).